Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

175. Proposed by M. E. GRABER, A. B., Instructor in Mathematics and Physics, Heidelberg University, Tiffin, Ohio.

Find the volume of the cono-cuneus determined by $z^2+a^2y^2/x^2=c^2$, which is contained between the planes $x=0$ and $x=a$. Ans. $\frac{1}{2}\pi c^2a$. [Todhunter's *Integral Calculus*, p. 189, No. 28.]

MECHANICS.

165. Proposed by O. W. ANTHONY, Head of the Mathematical Department, DeWitt Clinton High School, New York City.

Find the approximate form of a tower of circular cross section 1000 feet high and having a radius of lower base 20 feet, and so constructed that all the parts of the structure shall be subject to the same stress, due to the weight of the part of the tower above.

166. Proposed by G. B.M. ZERR, A. M., Ph. D., Parsons, West Va.

If a gravitating particle of mass m be placed at the point (a, b, c) prove that the work required to move a particle of unit mass from the point (x, y, z) to an infinite distance is $m[(x-a)^2+(y-b)^2+(z-c)^2]^{-\frac{1}{2}}$. Prove also that

$$\left(\frac{d^2}{dx^2}+\frac{d^2}{dy^2}+\frac{d^2}{dz^2}\right)\{m[(x-a)^2+(y-b)^2+(z-c)^2]^{-\frac{1}{2}}\}=0,$$

except when $x=a$, $y=b$, $z=c$.

MISCELLANEOUS.

143. Proposed by G. W. GREENWOOD, B. A. (Oxon), Professor of Mathematics and Astronomy, McKendree College, Lebanon, Ill.

If $A+B+C=180°$, show that $1-\cos^2A-\cos^2B-\cos^2C-2\cos A\cos B\cos C=0$.

144. Proposed by L. E. DICKSON. Ph. D., Assistant Professor of Mathematics, The University of Chicago.

Burnside, in his systematic search for all simple groups of orders from 661 to 1092 (*Proc. Lond. Math. Soc.*, 1895, pp. 333-338) overlooked the orders $792=2^3.3^2.11$ and $1008=2^4.3^2.7$. The former is immediately excluded. Discuss the latter order.

◆

NOTES.

Beginning with January, 1904, the *Mathematical Supplement of School Science* will be published bi-monthly, as a separate magazine entitled *School Mathematics*.

At the Christmas meeting of the San Francisco Section of the American Mathematical Society the following officers were elected for the coming year: Chairman, Professor R. E. Allardice; Secretary, Professor G. A. Miller; Program Committee, Professors M. W. Haskell, Irving Stringham, and G. A. Miller.